

## TLDR

Quick summary of today's notes. Lecture starts on next page.

- If  $T$  and  $U$  are functions  $\mathbb{R}^n \rightarrow \mathbb{R}^m$ , then there is a natural way to form the sum  $T + U$  and the scalar multiple  $cT$  for  $c \in \mathbb{R}$ . The definitions are

$$(T + U)(v) = T(v) + U(v) \quad \text{and} \quad (cT)(v) = c \cdot T(v) \quad \text{for } v \in \mathbb{R}^n.$$

Both of these are also functions  $\mathbb{R}^n \rightarrow \mathbb{R}^m$ .

If  $T$  and  $U$  are linear, then  $T + U$  and  $cT$  are both linear.

- If  $A$  and  $B$  are  $m \times n$  matrices, then there is a natural way to form the sum  $A + B$  and the scalar multiple  $cA$  for  $c \in \mathbb{R}$ . These operations work exactly the same as for vectors: we just add together entries in the same position or multiply all entries by the same number.

The resulting matrices have the same size as  $A$  and  $B$ .

- If  $A$  and  $B$  are the standard matrices of  $T$  and  $U$ , then  $A + B$  is the standard matrix of  $T + U$ , and  $cA$  for  $c \in \mathbb{R}$  is the standard matrix of  $cT$ .
- If  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $U : \mathbb{R}^k \rightarrow \mathbb{R}^n$  then we can *compose*  $T$  and  $U$  to form a new function

$$T \circ U : \mathbb{R}^k \rightarrow \mathbb{R}^m.$$

This function is defined by the formula  $T \circ U(v) = T(U(v))$  for  $v \in \mathbb{R}^k$ .

If  $T$  and  $U$  are linear then  $T \circ U$  is linear.

- There is a natural way to multiply an  $m \times n$  matrix  $A$  by an  $n \times k$  matrix  $B$ .

The result, written  $AB$ , is an  $m \times k$  matrix.

The product  $AB$  is only defined if the number of columns of  $A$  is the number of rows of  $B$ .

Unlike with scalars, we can have  $AB \neq BA$ , so the order of multiplication matters.

If  $A$  is the standard matrix of  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $B$  is the standard matrix of  $U : \mathbb{R}^k \rightarrow \mathbb{R}^n$ , then the product  $AB$  is the standard matrix of  $T \circ U : \mathbb{R}^k \rightarrow \mathbb{R}^m$ .

- To compute  $AB$ : if  $B = [b_1 \ b_2 \ \dots \ b_k]$  where  $b_i \in \mathbb{R}^n$  then  $AB = [Ab_1 \ Ab_2 \ \dots \ Ab_k]$ .

An example of this kind of calculation:

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 8 \end{bmatrix} \begin{bmatrix} 10 & 100 \\ 1 & 1000 \\ 10 & 100 \end{bmatrix} = \begin{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 8 \end{bmatrix} \begin{bmatrix} 10 \\ 1 \\ 10 \end{bmatrix} & \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 8 \end{bmatrix} \begin{bmatrix} 100 \\ 1000 \\ 100 \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 42 & 2400 \\ 84 & 4800 \end{bmatrix}.$$

- The transpose of a matrix  $A$  is the matrix  $A^T$  formed by flipping  $A$  across the diagonal, e.g.:

$$A = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} \quad \rightsquigarrow \quad A^T = \begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}.$$

If  $A$  is  $m \times n$  and  $B$  is  $n \times k$  so that the product  $AB$  is defined, then  $(AB)^T = B^T A^T$ .

# 1 Last time: one-to-one and onto linear transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a function.

The following mean the same thing:

- $T$  is *linear* in the sense that  $T(u + v) = T(u) + T(v)$  and  $T(cv) = cT(v)$  for all  $u, v \in \mathbb{R}^n$ ,  $c \in \mathbb{R}$ .
- There is an  $m \times n$  matrix  $A$  such that  $T$  has the formula  $T(v) = Av$  for all  $v \in \mathbb{R}^n$ .

If we are given a linear transformation  $T$ , then  $T(v) = Av$  for the matrix

$$A = [ T(e_1) \quad T(e_2) \quad \dots \quad T(e_n) ]$$

where  $e_i \in \mathbb{R}^n$  has a 1 in row  $i$  and 0 in all other rows. (If  $n = 2$  then  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .)

We call  $A$  the *standard matrix* of  $T$ .

The following all mean the same thing for a function  $f : X \rightarrow Y$ .

- $f$  is *one-to-one*.
- If  $a, b \in X$  and  $f(a) = f(b)$  then  $a = b$ .
- If  $a, b \in X$  and  $a \neq b$  then  $f(a) \neq f(b)$ .
- $f$  does not send different inputs to the same output.

Similarly, the following all mean the same thing for a function  $f : X \rightarrow Y$ .

- $f$  is *onto*.
- The range of  $f$  is equal to the codomain, i.e., the set  $\text{range}(f) = \{f(a) : a \in X\}$  is equal to  $Y$ .
- For each  $y \in Y$  there is at least one  $x \in X$  with  $f(x) = y$ .
- Every element of the codomain of  $f$  is an output for some input.

We can detect whether a linear transformation is one-to-one or onto by locating the pivot positions in its standard matrix (by row reducing).

**Theorem.** Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is the linear transformation  $T(v) = Av$  where  $A$  is an  $m \times n$  matrix.

- (1)  $T$  is one-to-one if and only if the columns of  $A$  are linearly independent, which happens precisely when  $A$  has a pivot position in every column.
- (2)  $T$  is onto if and only if the span of the columns of  $A$  is  $\mathbb{R}^m$ , which happens precisely when  $A$  has a pivot position in every row.

## 2 Operators on linear transformations and matrices

Key point from last time and starting point of today: linear transformations  $\mathbb{R}^n \rightarrow \mathbb{R}^m$  are uniquely represented by  $m \times n$  matrices, and every  $m \times n$  matrix corresponds to a linear transformation  $\mathbb{R}^n \rightarrow \mathbb{R}^m$ .

There are several simple, natural operations we can use to combine and alter linear transformations to get other linear transformations. The goal is to translate these function operations into matrix operations.

**Sums and scalar multiples.** Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $U : \mathbb{R}^n \rightarrow \mathbb{R}^m$  are two linear functions with the same domain and codomain. Their sum  $T + U$  is the function  $\mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by

$$(T + U)(v) = T(v) + U(v) \quad \text{for } v \in \mathbb{R}^n.$$

If  $c \in \mathbb{R}$  is a scalar, then  $cT$  is the function  $\mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by

$$(cT)(v) = cT(v) \quad \text{for } v \in \mathbb{R}^n.$$

**Fact.** Both  $T + U$  and  $cT$  are linear transformations.

*Proof.* To see that  $T + U$  is linear, we check that

$$(T + U)(u + v) = T(u + v) + U(u + v) = T(u) + T(v) + U(u) + U(v) = (T + U)(u) + (T + U)(v)$$

for  $u, v \in \mathbb{R}^n$ , and

$$(T + U)(av) = T(av) + U(av) = aT(v) + aU(v) = a(T + U)(v)$$

for  $a \in \mathbb{R}$  and  $v \in \mathbb{R}^n$ . Since these properties hold,  $T + U$  is linear.

The proof that  $cT$  is linear is similar. (Try this yourself!) □

Since sums and scalar multiples of linear functions are linear, it follows that differences  $T - U$  and arbitrary finite linear combinations  $aT + bU + cV + \dots$  of linear functions are linear.

Suppose  $T$  and  $U$  have standard matrices

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & & a_{2n} \\ \vdots & & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{22} & & b_{2n} \\ \vdots & & \ddots & \vdots \\ b_{m1} & b_{m2} & \dots & b_{mn} \end{bmatrix}$$

so that  $T(v) = Av$  and  $U(v) = Bv$  for all input vectors  $v \in \mathbb{R}^n$ .

**Proposition.** The standard matrix of  $T + U$  is the matrix  $A + B$  defined by

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \dots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & & a_{2n} + b_{2n} \\ \vdots & & \ddots & \vdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \dots & a_{mn} + b_{mn} \end{bmatrix}.$$

The standard matrix of  $cT$  is the matrix  $cA$  defined by

$$cA = \begin{bmatrix} ca_{11} & ca_{12} & \dots & ca_{1n} \\ ca_{21} & ca_{22} & & ca_{2n} \\ \vdots & & \ddots & \vdots \\ ca_{m1} & ca_{m2} & \dots & ca_{mn} \end{bmatrix}.$$

This is how we *define* sums and scalar multiples of matrices. These operations work in essentially the same way as for vectors: we can add matrices of the same size, by adding the entries in corresponding positions together, and we can multiply a matrix by a scalar  $c$  by multiplying all entries by  $c$ .

**Example.** We have

$$\begin{bmatrix} 4 & 0 & 5 \\ -1 & 3 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 3 & 5 & 7 \end{bmatrix} = \begin{bmatrix} 5 & 1 & 6 \\ 2 & 8 & 9 \end{bmatrix}.$$

and

$$-\begin{bmatrix} 4 & 0 & 5 \\ -1 & 3 & 2 \end{bmatrix} + 2\begin{bmatrix} 1 & 1 & 1 \\ 3 & 5 & 7 \end{bmatrix} = \begin{bmatrix} -4 & 0 & -5 \\ 1 & -3 & -2 \end{bmatrix} + \begin{bmatrix} 2 & 2 & 2 \\ 6 & 10 & 14 \end{bmatrix} = \begin{bmatrix} -2 & 2 & -3 \\ 7 & 7 & 12 \end{bmatrix}.$$

Suppose  $T, U, V$  are linear transformations  $\mathbb{R}^n \rightarrow \mathbb{R}^m$  with standard matrices  $A, B, C$ . Let  $a, b \in \mathbb{R}$ .

The following properties then hold:

<u>Functions</u>	<u>Matrices</u>
1. $T + U = U + T$	$A + B = B + A$ .
2. $(T + U) + V = T + (U + V)$	$(A + B) + C = A + (B + C)$ .
3. $T + 0 = T$ where $0 : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is the map $0(v) = 0 \in \mathbb{R}^m$ .	$A + 0 = A$ .
4. $a(T + U) = aT + aU$	$a(A + B) = aA + aB$ .
5. $(a + b)T = aT + bT$	$(a + b)A = aA + bA$ .
6. $a(bT) = (ab)T$ .	$a(bA) = (ab)A$ .

**Composition.** Suppose  $U : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $T : \mathbb{R}^m \rightarrow \mathbb{R}^k$  are functions.

Note that we assume the codomain of  $U$  is equal to the domain of  $T$ .

The *composition*  $T \circ U$  is the function  $\mathbb{R}^n \rightarrow \mathbb{R}^k$  given by

$$(T \circ U)(v) = T(U(v)) \quad \text{for } v \in \mathbb{R}^n.$$

**Fact.** Assume  $T$  and  $U$  are linear. Then  $T \circ U$  is linear.

*Proof.* To see that  $T \circ U$  is linear, we check that

$$(T \circ U)(u + v) = T(U(u + v)) = T(U(u) + U(v)) = T(U(u)) + T(U(v)) = (T \circ U)(u) + (T \circ U)(v)$$

for  $u, v \in \mathbb{R}^n$ , and

$$(T \circ U)(cv) = T(U(cv)) = T(cU(v)) = cT(U(v)) = c(T \circ U)(v)$$

for  $c \in \mathbb{R}$  and  $v \in \mathbb{R}^n$ . □

Important note:  $U \circ T$  is not defined unless  $k = n$ .

Even if  $k = n$  so that both  $T \circ U$  and  $U \circ T$  are defined, there is no reason to expect that  $T \circ U = U \circ T$ .

**Example.** If  $n = m = k = 1$  and  $T(x) = 2x$  and  $U(x) = x^2$ , then

$$(T \circ U)(x) = T(x^2) = 2x^2 \quad \text{but} \quad (U \circ T)(x) = U(2x) = 4x^2.$$

Assume  $U : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $T : \mathbb{R}^m \rightarrow \mathbb{R}^k$  are linear.

Then  $T \circ U$  is a linear transformation  $\mathbb{R}^n \rightarrow \mathbb{R}^k$ , so there is a unique  $k \times n$  matrix  $C$  such that

$$(T \circ U)(v) = Cv \quad \text{for } v \in \mathbb{R}^n.$$

If  $A$  is the standard matrix of  $T$  and  $B$  is the standard matrix of  $U$ , then we define the *matrix product*

$$AB = C.$$

Note how this definition works: if  $A$  is  $k \times m$  and  $B$  is  $m \times n$  then we *define*  $AB$  to be the unique  $k \times n$  matrix  $C$  such that  $Cv = A(Bv)$  for all  $v \in \mathbb{R}^n$ .

How do we actually compute the rectangular array  $AB$  from  $A$  and  $B$ ?

**Theorem.** Suppose  $B$  has columns  $b_1, b_2, \dots, b_n \in \mathbb{R}^m$  so that  $B = [ b_1 \ b_2 \ \dots \ b_n ]$ .

Then  $AB = [ Ab_1 \ Ab_2 \ \dots \ Ab_n ]$ . (This makes sense as  $A$  is  $k \times m$ .)

*Proof.*  $AB$  is the standard matrix of the linear function  $T \circ U$ , so

$$\begin{aligned} AB &= [ (T \circ U)(e_1) \ (T \circ U)(e_2) \ \dots \ (T \circ U)(e_n) ] \\ &= [ T(U(e_1)) \ T(U(e_2)) \ \dots \ T(U(e_n)) ] \\ &= [ A(Be_1) \ A(Be_2) \ \dots \ A(Be_n) ] \\ &= [ Ab_1 \ Ab_2 \ \dots \ Ab_n ]. \end{aligned}$$

□

**Example.** If  $A = \begin{bmatrix} 2 & 3 \\ 1 & -5 \end{bmatrix}$  and  $B = \begin{bmatrix} 4 & 3 & 6 \\ 1 & -2 & 3 \end{bmatrix}$ , then  $b_1 = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$ ,  $b_2 = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$ ,  $b_3 = \begin{bmatrix} 6 \\ 3 \end{bmatrix}$ , so

$$AB = [ Ab_1 \ Ab_2 \ Ab_3 ] = \begin{bmatrix} 11 & 0 & 21 \\ -1 & 13 & -9 \end{bmatrix}.$$

The quick rule for computing  $AB$ : if the  $i$ th row of  $A$  and  $j$ th column of  $B$  are

$$\begin{bmatrix} a_1 & a_2 & \dots & a_m \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

then the entry in the  $i$ th row and  $j$ th column of  $AB$  is

$$\begin{bmatrix} a_1 & a_2 & \dots & a_m \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} = a_1 b_1 + a_2 b_2 + \dots + a_m b_m.$$

**Example.** Suppose  $A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \\ 9 & 9 & 9 \end{bmatrix}$ .

The entry in the 2nd row and 2nd column of  $AB$  is

$$5 \cdot 2 + 6 \cdot 5 + 7 \cdot 8 + 8 \cdot 9 = 10 + 30 + 56 + 72 = 168.$$

A position  $(i, j)$  in a matrix is *diagonal* if  $i = j$ .

Write  $I_n$  for the  $n \times n$  matrix

$$I_n = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{bmatrix}$$

which has 1 in each diagonal position, and zeros in all other positions.

The matrix  $I_n$  is the standard matrix of the *identity map*  $\mathbb{R}^n \rightarrow \mathbb{R}^n$ .

This is the linear function  $T$  with  $T(v) = v$  for all  $v \in \mathbb{R}^n$ .

**Proposition.** Let  $A, B, C$  be matrices.

Assume  $A$  is  $m \times n$ ,  $B$  is  $n \times l$ , and  $C$  is  $l \times k$ .

Then  $A(BC) = (AB)C$ .

*Proof.* Suppose  $A, B$ , and  $C$  are the standard matrices of linear transformations  $T_A, T_B, T_C$ .

It holds that  $T_A \circ (T_B \circ T_C) = (T_A \circ T_B) \circ T_C$ , since for any input  $x$  we have

$$(T_A \circ (T_B \circ T_C))(x) = T_A((T_B \circ T_C)(x)) = T_A(T_B(T_C(x)))$$

which is the same thing as

$$((T_A \circ T_B) \circ T_C)(x) = (T_A \circ T_B)(T_C(x)) = T_A(T_B(T_C(x))).$$

In fact, this holds for any functions  $T_A, T_B$ , and  $T_C$  that we can compose together; nothing here depends on the fact that these are *linear* functions.

However, since  $T_A, T_B, T_C$  are linear, both  $T_A \circ (T_B \circ T_C)$  and  $(T_A \circ T_B) \circ T_C$  are linear, and they have the same standard matrix since they are equal as functions.

As  $A(BC)$  is the standard matrix of  $T_A \circ (T_B \circ T_C)$ , while  $(AB)C$  is the standard matrix of  $(T_A \circ T_B) \circ T_C$ , we must have  $A(BC) = (AB)C$ .  $\square$

Here are some easier properties. Suppose  $A, B, C$  are matrices and  $r \in \mathbb{R}$ .

- If  $A$  is  $m \times n$  and  $B, C$  are  $n \times l$  then  $A(B + C) = AB + AC$ .
- If  $A, B$  are  $m \times n$  and  $C$  is  $n \times l$  then  $(A + B)C = AC + BC$ .
- If  $A$  is  $m \times n$  and  $B$  is  $n \times l$  then  $r(AB) = (rA)B = A(rB)$ .
- If  $A$  is  $m \times n$  then  $I_m A = A I_n = A$ .

### 3 Pathologies of matrix multiplication

Suppose  $A$  and  $B$  are matrices.

Four important observations:

1. The product  $AB$  is defined only if the number of columns of  $A$  is the number of rows of  $B$ .
2. Even if  $AB$  and  $BA$  are both defined, it often happens that  $AB \neq BA$ .
3.  $AB = AC$  does not imply  $B = C$ , and  $AC = BC$  does not imply  $A = B$ .

4. It can happen that  $AB = 0 = \begin{bmatrix} 0 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 0 \end{bmatrix}$  even if both  $A \neq 0$  and  $B \neq 0$ .

**Example.** We have

$$\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} & \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

while

$$\begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} & \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 2 & 0 \end{bmatrix}.$$

If  $A$  and  $B$  are both *square* matrices of the same size (meaning they have the same number of rows and columns), and  $AB = BA$ , then we say that  $A$  and  $B$  *commute*.

## 4 Matrix transpose

The *transpose* of an  $m \times n$  matrix  $A$  is the  $n \times m$  matrix  $A^T$  whose columns are the rows of  $A$ .

If  $a_{ij}$  is the entry in row  $i$  and column  $j$  of  $A$ , then this is the entry in row  $j$  and column  $i$  of  $A^T$ .

For example, if  $A = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix}$  then  $A^T = \begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}$ .

The transpose of  $A$  is given by flipping  $A$  across the main diagonal, in order to interchange rows/columns.

Another example: if  $C = \begin{bmatrix} 1 & 1 & 1 & 1 \\ -3 & 5 & -2 & 7 \\ 0 & 0 & 1 & 0 \end{bmatrix}$  then  $C^T = \begin{bmatrix} 1 & -3 & 0 \\ 1 & 5 & 0 \\ 1 & -2 & 1 \\ 1 & 7 & 0 \end{bmatrix}$ .

We finish this lecture by noting some basic properties of the transpose operation:

- $(A^T)^T = A$  since flipping twice does nothing.
- If  $A$  and  $B$  have the same size then  $(A + B)^T = A^T + B^T$ .
- If  $c \in \mathbb{R}$  then  $(cA)^T = c(A^T)$ .
- If  $A$  is an  $k \times m$  matrix and  $B$  is an  $m \times n$  matrix then  $(AB)^T = B^T A^T$ .

To prove the last property, use our earlier results to compute the entries in  $i$ th row and  $j$ th column of the matrices on either side (in terms of the entries of  $A$  and  $B$ ), and check that these are equal.

Questions for later: how is the linear transformation with standard matrix  $A$  related to the linear transformation with standard matrix  $A^T$ ? What is the transpose as an operation on linear transformations?

## 5 Vocabulary

Keywords from today's lecture:

1. **Sums, scalar multiples, and compositions** of linear functions.

If  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $U : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $c \in \mathbb{R}$  then

$$T + U : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

is the function with  $(T + U)(v) = T(v) + U(v)$ , and

$$cT : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

is the function with  $(cT)(v) = c(T(v))$ .

If  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $U : \mathbb{R}^m \rightarrow \mathbb{R}^k$  then  $U \circ T : \mathbb{R}^n \rightarrow \mathbb{R}^k$  is the function  $(U \circ T)(v) = U(T(v))$ .

2. **Sums, scalar multiples, and products** of matrices.

If  $A$  and  $B$  are  $m \times n$  matrices then  $A + B$  is the  $m \times n$  matrix whose entry in position  $(i, j)$  is  $A_{ij} + B_{ij}$ . If  $c \in \mathbb{R}$  then  $cA$  is the matrix whose entry in position  $(i, j)$  is  $cA_{ij}$ .

If  $A$  is  $m \times n$  and  $B$  is  $n \times k$  then  $AB$  is the  $m \times k$  whose entry in position  $(i, j)$  is the  $i$ th row of  $A$  (which is a  $1 \times n$  matrix) times the  $j$ th column of  $B$  (which is a vector in  $\mathbb{R}^n$ ).

Example: 
$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} + \begin{bmatrix} w & x \\ y & z \end{bmatrix} = \begin{bmatrix} a+w & b+x \\ c+y & d+z \end{bmatrix}.$$

Example: 
$$5 \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 5a & 5b \\ 5c & 5d \end{bmatrix}.$$

Example: 
$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} w & x \\ y & z \end{bmatrix} = \begin{bmatrix} aw+by & ax+bz \\ cw+dy & cx+dz \end{bmatrix}.$$

3. **Transpose** of a matrix.

If  $A$  is an  $m \times n$  matrix then its transpose  $A^T$  is the  $n \times m$  with the same entries as  $A$  but with rows and columns interchanged.

Example: 
$$\begin{bmatrix} a & b & c \\ x & y & z \end{bmatrix}^T = \begin{bmatrix} a & x \\ b & y \\ c & z \end{bmatrix}.$$

4. **Identity** matrix

The  $n \times n$  matrix  $I = I_n$  with 1s on the diagonal and 0s off the diagonal.

$AI = A$  and  $IB = B$  for all matrices  $A$  with  $n$  columns and all matrices  $B$  with  $n$  rows.

Example:  $[1]$ ,  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ,  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ , and so on.