

TLDR

Quick summary of today's notes. Lecture starts on next page.

- A *line of best fit* through data points $(a_1, b_1), (a_2, b_2), \dots, (a_n, b_n)$ is an equation of the form

$$y = \beta_0 + \beta_1 x$$

where $\begin{bmatrix} \beta_0 \\ \beta_1 \end{bmatrix} \in \mathbb{R}^2$ is a least-squares solution to $Ax = b$ where $A = \begin{bmatrix} 1 & a_1 \\ 1 & a_2 \\ \vdots & \vdots \\ 1 & a_n \end{bmatrix}$ and $b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$.

- A matrix A is *symmetric* if $A^T = A$. This can only hold if A is square. For example:

$$\begin{bmatrix} 0 & -1 & 0 \\ -1 & 5 & 8 \\ 0 & 8 & -7 \end{bmatrix}.$$

If A is symmetric then so is A^2, A^3, A^4 , etc.

If A is symmetric and invertible then so is A^{-1}, A^{-2}, A^{-3} , etc.

If A is symmetric and u and v are eigenvectors for A with *different* eigenvalues, then $u \bullet v = 0$.

- A list of vectors u_1, u_2, \dots, u_p is *orthonormal* if $u_i \bullet u_i = 1$ and $u_i \bullet u_j = 0$ for all $i \neq j$.

A square matrix P is invertible with $P^{-1} = P^T$ if and only if its columns are orthonormal.

An $n \times n$ matrix A is *orthogonally diagonalizable* if there is a diagonal matrix D and an invertible matrix P with $P^{-1} = P^T$ such that $A = PDP^{-1}$.

- When we have such a decomposition $A = PDP^{-1}$ where D is diagonal and $P^{-1} = P^T$, the diagonal entries of D are the eigenvalues of A , and the columns of P are an orthonormal basis of \mathbb{R}^n consisting of eigenvectors for A .

Conversely, an $n \times n$ matrix A is orthogonally diagonalizable if and only if there exists an orthonormal basis of \mathbb{R}^n consisting of eigenvectors for A .

- Surprising fact: all (complex) eigenvalues of a symmetric matrix belong to \mathbb{R} .

Surprising fact: an $n \times n$ matrix A is orthogonally diagonalizable if and only if $A = A^T$.

Much of this lecture is spent proving these facts.

- To orthogonally diagonalize a given $n \times n$ symmetric matrix A , you need to find an orthogonal basis of \mathbb{R}^n consisting of eigenvectors v_1, v_2, \dots, v_n for A .

Once you find this, let $u_i = \frac{1}{\|v_i\|} v_i$ and $U = \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix}$.

Then $A = UDU^T$ where D is the diagonal matrix whose i th diagonal entry is the eigenvalue of v_i .

To find the orthogonal basis of eigenvectors v_1, v_2, \dots, v_n :

1. Factor the characteristic polynomial of A to compute its eigenvalues.
2. For each eigenvalue λ , do the usual row reduce procedure to find a basis for $\text{Nul}(A - \lambda I)$.
3. Apply the Gram-Schmidt process to convert your basis of $\text{Nul}(A - \lambda I)$ to an orthogonal basis.
4. Finally combine these orthogonal bases — the combined list of vectors will still be orthogonal.

1 Last time: least-squares problems

Definition. Suppose A is an $m \times n$ matrix and $b \in \mathbb{R}^m$.

The linear system $A^T Ax = A^T b$ is always consistent, i.e., has at least one solution.

A solution to $A^T Ax = A^T b$ is called a *least-squares solution* to $Ax = b$.

Let $\|v\| = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2} \geq 0$ for $v \in \mathbb{R}^n$. Recall that $\|v\| = 0$ if and only if $v = 0$.

Fact. A vector $s \in \mathbb{R}^n$ is a least-squares solution to $Ax = b$ if and only if $\|b - As\| \leq \|b - Ax\|$ for all x .

The linear system $Ax = b$ is consistent if and only if $\|b - Ax\| = 0$ for some $x \in \mathbb{R}^n$.

This means that if $Ax = b$ is consistent then all least-squares solutions s satisfy $\|b - As\| = 0$ so $As = b$.

If $Ax = b$ is inconsistent, there is still at least one least-squares solution s (but in this case $\|b - As\| > 0$).

Theorem. Let A be an $m \times n$ matrix. The following properties are equivalent:

- (a) $Ax = b$ has a unique least-squares solution for each $b \in \mathbb{R}^m$.
- (b) The columns of A are linearly independent.
- (c) $A^T A$ is invertible.

Example (Lines of best fit). Suppose we have n data points $(a_1, b_1), (a_2, b_2), \dots, (a_n, b_n)$.

We want to find parameters $\beta_0, \beta_1 \in \mathbb{R}$ such that $y = \beta_0 + \beta_1 x$ describes the *line of best fit* for this data.

If our points are all on the same line, then for some $\begin{bmatrix} \beta_0 \\ \beta_1 \end{bmatrix} \in \mathbb{R}^2$ we would have

$$b_i = \beta_0 + \beta_1 a_i \quad \text{for } i = 1, 2, \dots, n,$$

meaning that $x = \begin{bmatrix} \beta_0 \\ \beta_1 \end{bmatrix}$ is an exact solution to the linear system $Ax = b$ where

$$A = \begin{bmatrix} 1 & a_1 \\ 1 & a_2 \\ \vdots & \vdots \\ 1 & a_n \end{bmatrix} \quad \text{and} \quad b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}.$$

If the given points are not on the same line, then no exact solution to $Ax = b$ exists, and we should instead try to find a least-squares solution to this linear system.

To be concrete, suppose we have four points $(2, 1), (5, 2), (7, 3)$, and $(8, 3)$ so that

$$A = \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \quad \text{and} \quad b = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

The least-squares solutions to $Ax = b$ are the exact solutions to $A^T Ax = A^T b$. We have

$$A^T A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$


and

$$A^T b = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}.$$

The matrix $A^T A$ is invertible. (Why?) It follows that a least-squares solution is provided by

$$\begin{bmatrix} \beta_0 \\ \beta_1 \end{bmatrix} = (A^T A)^{-1} A^T b = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}^{-1} \begin{bmatrix} 9 \\ 57 \end{bmatrix} = \frac{1}{84} \begin{bmatrix} 142 & -22 \\ -22 & 4 \end{bmatrix} \begin{bmatrix} 9 \\ 57 \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}.$$

Thus our line of best fit for the data is $y = \frac{2}{7} + \frac{5}{14}x$:



2 Symmetric matrices

A matrix A is *symmetric* if $A^T = A$. This happens if A is square and $A_{ij} = A_{ji}$ for all i, j .

Example. $\begin{bmatrix} 1 & 0 \\ 0 & -3 \end{bmatrix}$ and $\begin{bmatrix} 0 & -1 & 0 \\ -1 & 5 & 8 \\ 0 & 8 & -7 \end{bmatrix}$ and $\begin{bmatrix} a & b & c \\ b & d & e \\ c & e & f \end{bmatrix}$ are symmetric matrices.

$\begin{bmatrix} 1 & -3 \\ 3 & 0 \end{bmatrix}$ and $\begin{bmatrix} 1 & -4 & 0 \\ -6 & 1 & -4 \\ 6 & -6 & 1 \end{bmatrix}$ and $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 5 \end{bmatrix}$ are not symmetric.

Proposition. If A is a symmetric matrix and k is a positive integer then A^k is also symmetric.

Proof. If $A = A^T$ then $(A^k)^T = (AA \cdots A)^T = A^T \cdots A^T A^T = (A^T)^k = A^k$. □

Proposition. If A is an invertible symmetric matrix then A^{-1} is also symmetric.

Proof. This is because $(A^{-1})^T = (A^T)^{-1}$. □

Recall how we can diagonalize a matrix.

Example. Let $A = \begin{bmatrix} 6 & -2 & -1 \\ -2 & 6 & -1 \\ -1 & -1 & 5 \end{bmatrix}$.

Then $\det(A - xI) = (8 - x)(6 - x)(3 - x)$ so the eigenvalues of A are 8, 6, and 3. By constructing bases for the null spaces of $A - 8I$, $A - 6I$, and $A - 3I$, we find that the following are eigenvectors of A :

$$v_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} \text{ with eigenvalue } 8.$$

$$v_2 = \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix} \text{ with eigenvalue } 6.$$

$$v_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \text{ with eigenvalue } 3.$$

These eigenvectors are actually an orthogonal basis for \mathbb{R}^3 .

Converting these vectors to unit vectors gives an orthonormal basis of eigenvectors:

$$u_1 = \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix}, \quad u_2 = \begin{bmatrix} -1/\sqrt{6} \\ -1/\sqrt{6} \\ 2/\sqrt{6} \end{bmatrix}, \quad u_3 = \begin{bmatrix} 1/\sqrt{3} \\ 1/\sqrt{3} \\ 1/\sqrt{3} \end{bmatrix}.$$

We then have $A = PDP^{-1}$ where

$$P = [u_1 \quad u_2 \quad u_3] \quad \text{and} \quad D = \begin{bmatrix} 8 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 3 \end{bmatrix}.$$

(Why does this hold? It is enough to check that $PDP^{-1}v = Av$ for $v \in \{u_1, u_2, u_3\}$.)

Since the columns of P are orthonormal, we actually have $P^T = P^{-1}$ so $A = PDP^T$.

The special properties in this example will turn out to hold for all symmetric matrices.

Theorem. Suppose A is a symmetric matrix. Then any two eigenvectors from different eigenspaces of A are orthogonal. In other words, if $A = A^T$ is $n \times n$ and $u, v \in \mathbb{R}^n$ are such that $Au = au$ and $Av = bv$ for numbers $a, b \in \mathbb{R}$ with $a \neq b$, then $u \bullet v = 0$.

Proof. Let u and v be eigenvectors of A with eigenvalues a and b , where $a \neq b$.

Then $au \bullet v = Au \bullet v = (Au)^T v = u^T A^T v = u^T Av = u \bullet Av = u \bullet bv$.

But $au \bullet v = a(u \bullet v)$ and $u \bullet bv = b(u \bullet v)$, so this means $a(u \bullet v) = b(u \bullet v)$ and therefore $(a - b)(u \bullet v) = 0$.

Since $a - b \neq 0$, it follows that $u \bullet v = 0$. □

Recall that a matrix P is *orthogonal* if P is invertible and $P^{-1} = P^T$.

Definition. A matrix A is *orthogonally diagonalizable* if there is an orthogonal matrix P and a diagonal matrix D such that $A = PDP^{-1} = PDP^T$.

When A is orthogonally diagonalizable and $A = PDP^{-1} = PDP^T$, the diagonal entries of D are the eigenvalues of A , and the columns of P are the corresponding eigenvectors; moreover, these eigenvectors form an orthonormal basis of \mathbb{R}^n .

In fact, it follows by the arguments in our earlier lectures about diagonalizable matrices that an $n \times n$ matrix A is orthogonally diagonalizable if and only if there is an orthonormal basis for \mathbb{R}^n consisting of eigenvectors for A .

Surprisingly, there is a much more direct characterization of orthogonally diagonalizable matrices:

Theorem. A square matrix is orthogonally diagonalizable if and only if it is symmetric.

We prove this after a sequence of lemmas.

Lemma. If A is orthogonally diagonalizable then A is symmetric.

Proof. If X, Y, Z are $n \times n$ matrices then $(XYZ)^T = Z^T(XY)^T = Z^TY^TX^T$.

Suppose $A = PDP^T$ where D is diagonal. Then $D = D^T$ and $(P^T)^T = P$, so

$$A^T = (PDP^T)^T = (P^T)^T D^T P^T = PDP^T = A.$$

□

Lemma. All (complex) eigenvalues of an $n \times n$ symmetric matrix A with real entries belong to \mathbb{R} .

Proof. Suppose A is a symmetric $n \times n$ matrix with real entries, so that $A = A^T = \overline{A}$.

Let $v \in \mathbb{C}^n$. Then $\overline{v}^T Av$ is some complex number.

For example, if $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $v = \begin{bmatrix} 1+i \\ 1-i \end{bmatrix}$ then

$$\overline{v}^T Av = \begin{bmatrix} 1-i & 1+i \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1+i \\ 1-i \end{bmatrix} = \begin{bmatrix} 3+i & 3-i \end{bmatrix} \begin{bmatrix} 1+i \\ 1-i \end{bmatrix} = (3+i)(1+i) + (3-i)(1-i) = 4.$$

In fact, the number $\overline{v}^T Av$ belongs to \mathbb{R} since $\overline{\overline{v}^T Av} = v^T A \overline{v} = (\overline{v}^T Av)^T = \overline{v}^T Av$.

(The last equality holds since both sides are 1×1 matrices, i.e., scalars.)

Now suppose $v \in \mathbb{C}^n$ is an eigenvector for A with eigenvalue $\lambda \in \mathbb{C}$. Then $\overline{v}^T Av = \overline{v}^T (\lambda v) = \lambda (\overline{v}^T v) \in \mathbb{R}$. The complex number $\overline{v}^T v$ always belongs to \mathbb{R} (why?) so it must also hold that $\lambda \in \mathbb{R}$. □

Lemma. An $n \times n$ matrix A with all real eigenvalues can be written as $A = URU^T$ where U is an $n \times n$ orthogonal matrix (i.e., has orthonormal columns) and R is an $n \times n$ upper-triangular matrix.

One calls $A = URU^T$ with U and R of this form a *Schur factorization* of A .

Proof. Suppose A is an $n \times n$ matrix with all real eigenvalues.

Let $u_1 \in \mathbb{R}^n$ be a unit eigenvector for A with eigenvalue $\lambda \in \mathbb{R}$.

Let $u_2, \dots, u_n \in \mathbb{R}^n$ be any vectors such that u_1, u_2, \dots, u_n is an orthonormal basis for \mathbb{R}^n .

(One way to construct these vectors: let $u_1 = x_1, x_2, \dots, x_n$ be any basis, apply the Gram-Schmidt process to get $u_1 = v_1, v_2, \dots, v_n$, and then convert each v_i to a unit vector.)

Define $U = \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix}$ so that $U^T = U^{-1}$.

By considering the product $U^T AU e_i$ for $i = 1, 2, \dots, n$, one finds that $U^T AU$ has the form

$$U^T AU = \begin{bmatrix} \lambda & * \\ 0 & B \end{bmatrix}$$

for some $(n-1) \times (n-1)$ matrix B . Here, $*$ stands for $n-1$ arbitrary entries.

The matrix $U^T AU = U^{-1}AU$ has the same characteristic polynomial as A .

This polynomial is just $(\lambda - x) \det(B - xI)$, which is $\lambda - x$ times the characteristic polynomial of B .

Since the characteristic polynomial of A has all real roots, the same must be true of the characteristic polynomial of B . Thus B must also have all real eigenvalues.

By repeating the argument above, we deduce that there is an eigenvalue $\mu \in \mathbb{R}$ for B , an $(n-1) \times (n-1)$ orthogonal matrix V , and an $(n-2) \times (n-2)$ matrix C with all real eigenvalues such that

$$V^T BV = \begin{bmatrix} \mu & * \\ 0 & C \end{bmatrix}.$$

The matrix $\begin{bmatrix} 1 & 0 \\ 0 & V \end{bmatrix}$ is also orthogonal, and the product of orthogonal matrices is orthogonal. (Why?)

It follows for the orthogonal matrix $W = U \begin{bmatrix} 1 & 0 \\ 0 & V \end{bmatrix}$ that $W^T A W = \begin{bmatrix} \lambda & * & * \\ 0 & \mu & * \\ 0 & 0 & C \end{bmatrix}$.

By continuing in this way, we will eventually construct an orthogonal matrix X and an upper-triangular matrix R such that $X^T A X = R$, in which case $A = X X^T A X X^T = X R X^T$. \square

Now we can prove the theorem.

Proof of theorem. The first lemma shows that if A is orthogonally diagonalizable then A is symmetric.

Suppose conversely that A is symmetric. Then A has all real eigenvalues, so there exists a Schur factorization $A = U R U^T$. We then have $A^T = (U R U^T)^T = U R^T U^T$ but also $A^T = A = U R U^T$.

Since $U^T = U^{-1}$, it follows that $R = R^T$. Since R is upper-triangular, this can only hold if R is diagonal.

But if R is diagonal then $A = U R U^T$ is orthogonally diagonalizable. \square

To orthogonally diagonalize an $n \times n$ symmetric matrix A , we just need to find an orthogonal basis of eigenvectors v_1, v_2, \dots, v_n for \mathbb{R}^n . Then $A = U D U^T$ with $U = [u_1 \ u_2 \ \dots \ u_n]$ where $u_i = \frac{1}{\|v_i\|} v_i$ and D is the diagonal matrix of the corresponding eigenvalues.

If all eigenspaces of A are 1-dimensional, then any basis of eigenvectors will be orthogonal. If A has an eigenspace of dimension greater than one, then after finding a basis for this eigenspace, it is necessary to apply the Gram-Schmidt process to convert this basis to one that is orthogonal.

Corollary. If $A = U D U^T$ where $U = [u_1 \ u_2 \ \dots \ u_n]$ has orthonormal columns and

$$D = \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{bmatrix}$$

is diagonal, then $A = \lambda_1 u_1 u_1^T + \lambda_2 u_2 u_2^T + \dots + \lambda_n u_n u_n^T$.

Each product $u_i u_i^T$ is an $n \times n$ matrix of rank 1. One calls this expression a *spectral decomposition* of A .

Example. Let $A = \begin{bmatrix} 7 & 2 \\ 2 & 4 \end{bmatrix}$. A spectral decomposition of A is given by

$$\begin{aligned} A &= \begin{bmatrix} 2/\sqrt{5} & -1/\sqrt{5} \\ 1/\sqrt{5} & 2/\sqrt{5} \end{bmatrix} \begin{bmatrix} 8 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 2/\sqrt{5} & 1/\sqrt{5} \\ -1/\sqrt{5} & 2/\sqrt{5} \end{bmatrix} \\ &= 8 \begin{bmatrix} 2/\sqrt{5} \\ 1/\sqrt{5} \end{bmatrix} \begin{bmatrix} 2/\sqrt{5} & 1/\sqrt{5} \end{bmatrix} + 3 \begin{bmatrix} -1/\sqrt{5} \\ 2/\sqrt{5} \end{bmatrix} \begin{bmatrix} -1/\sqrt{5} & 2/\sqrt{5} \end{bmatrix} \\ &= \begin{bmatrix} 32/5 & 16/5 \\ 16/5 & 8/5 \end{bmatrix} + \begin{bmatrix} 3/5 & -6/5 \\ -6/5 & 12/5 \end{bmatrix}. \end{aligned}$$

3 Vocabulary

Keywords from today's lecture:

1. *Symmetric matrix.*

A matrix A that is equal to its transpose, so that $A = A^T$. Such a matrix is square.

Symmetric matrices are precisely the square matrices A that are *orthogonally diagonalizable*, in other words, the matrices that can be expressed as

$$A = PDP^T$$

where D is a diagonal matrix and P is an invertible matrix with $P^{-1} = P^T$.

Example: $\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$ or any diagonal matrix.

2. *Schur factorization* of an $n \times n$ matrix A .

A decomposition $A = URU^T$ where R is an $n \times n$ upper triangular matrix and U is an orthogonal matrix (i.e., U is invertible with $U^{-1} = U^T$).