

MATH2013 Multivariable Calculus

From the textbook Calculus - Several Variables (5th) by R. Adams, Addison/Wesley/Longman.

* **At least try to do the underlined ones, the others are recommended exercises.**

Homework 1

(Total: 21 questions)

Ex. 10.1

- 6 Show that the triangle with vertices $(1, 2, 3)$, $(4, 0, 5)$, and $(3, 6, 4)$ has a right angle.
- 14 Describe (and sketch if possible) the set of points in \mathbb{R}^3 which satisfy the equation $z = x$.
- 22 Describe (and sketch if possible) the set of points in \mathbb{R}^3 which satisfy the inequality $z \geq \sqrt{x^2 + y^2}$.

Ex. 10.2

- 2 Calculate the following for the given vectors \mathbf{u} and \mathbf{v} :

$$\mathbf{u} = \mathbf{i} - \mathbf{j} \quad \text{and} \quad \mathbf{v} = \mathbf{j} + 2\mathbf{k}.$$

- (a) $\mathbf{u} + \mathbf{v}$, $\mathbf{u} - \mathbf{v}$, $2\mathbf{u} - 3\mathbf{v}$,
- (b) the lengths $\|\mathbf{u}\|$ and $\|\mathbf{v}\|$,
- (c) unit vectors $\hat{\mathbf{u}}$ and $\hat{\mathbf{v}}$ in the directions of \mathbf{u} and \mathbf{v} , respectively,
- (d) the dot product $\mathbf{u} \cdot \mathbf{v}$,
- (e) the angle between \mathbf{u} and \mathbf{v} ,
- (f) the scalar projection of \mathbf{u} in the direction of \mathbf{v} ,
- (g) the vector projection of \mathbf{v} along \mathbf{u} .
- 10 A straight river 500m wide flows due east at a constant speed of 3 km/h. If you can row your boat at speed of 5 km/h in still water, in what direction should you head if you wish to row from point A on the south shore to point B on the north shore directly north of A ? How long will the trip take?

- 16 If a vector \mathbf{u} in \mathbb{R}^3 makes angles α , β , and γ with the coordinate axes, show that

$$\hat{\mathbf{u}} = \cos \alpha \mathbf{i} + \cos \beta \mathbf{j} + \cos \gamma \mathbf{k}$$

is a unit vector in the direction of \mathbf{u} . Hence show that

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1.$$

- 19 If \mathbf{r}_1 and \mathbf{r}_2 are the position vectors of two points, P_1 and P_2 , and λ is a real number, show that

$$\mathbf{r} = (1 - \lambda)\mathbf{r}_1 + \lambda\mathbf{r}_2$$

is the position vector of a point P on the straight line joining P_1 and P_2 . Where is P if $\lambda = 1/2$? if $\lambda = 2/3$? if $\lambda = -1$? if $\lambda = 2$?

- 20 Let \mathbf{a} be a nonzero vector. Describe the set of all points in 3-space whose position vectors \mathbf{r} satisfy $\mathbf{a} \cdot \mathbf{r} = 0$.
- 24 Let $\mathbf{u} = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k}$, $\mathbf{v} = \mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$, and $\mathbf{w} = 2\mathbf{i} - 2\mathbf{j} + \mathbf{k}$. Find two unit vectors each of which makes equal angles with \mathbf{u} , \mathbf{v} , and \mathbf{w} .
- 27 Let \mathbf{u} and \mathbf{v} be two vectors.
- (a) Show that $\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + 2\mathbf{u} \cdot \mathbf{v} + \|\mathbf{v}\|^2$.
- (b) Show that $\mathbf{u} \cdot \mathbf{v} \leq \|\mathbf{u}\| \|\mathbf{v}\|$.
- (c) Deduce from (a) and (b) that $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$.

- 29 Let $\mathbf{u} = \frac{3}{5}\mathbf{i} + \frac{4}{5}\mathbf{j}$, $\mathbf{v} = \frac{4}{5}\mathbf{i} - \frac{3}{5}\mathbf{j}$, and $\mathbf{w} = \mathbf{k}$.

- (a) Show that $\|\mathbf{u}\| = \|\mathbf{v}\| = \|\mathbf{w}\| = 1$ and $\mathbf{u} \cdot \mathbf{v} = \mathbf{u} \cdot \mathbf{w} = 0$. The vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} are mutually perpendicular unit vectors, and as such are said to constitute an **orthonormal basis** for \mathbb{R}^3 .
- (b) If $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, show by direct calculation that

$$\mathbf{r} = (\mathbf{r} \cdot \mathbf{u})\mathbf{u} + (\mathbf{r} \cdot \mathbf{v})\mathbf{v} + (\mathbf{r} \cdot \mathbf{w})\mathbf{w}.$$

- 33 Given constants r , s and t , with $r \neq 0$ and $s \neq 0$, and given a vector \mathbf{a} satisfying $\|\mathbf{a}\|^2 > 4rst$, solve the system of equations

$$r\mathbf{x} + s\mathbf{y} = \mathbf{a}$$

$$\mathbf{x} \cdot \mathbf{y} = t$$

for the unknown vectors \mathbf{x} and \mathbf{y} .

Ex. 10.3

- 14 (**Volume of a tetrahedron**) A **tetrahedron** is a pyramid with a triangular base and three other triangular faces. It has four vertices and six edges. Like any pyramid or cone, its volume is equal to $\frac{1}{3}AH$, where A is the area of the base and H is the height measured perpendicular

to the base. If \mathbf{u} , \mathbf{v} and \mathbf{w} are vectors coinciding with the three edges of a tetrahedron that meet at one vertex, show that the tetrahedron has volume given by

$$\text{Volume} = \frac{1}{6} \|\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})\| = \frac{1}{6} \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix}.$$

20 If $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = 0$ but $\mathbf{v} \times \mathbf{w} \neq \mathbf{0}$, show that there are constants λ and μ such that

$$\mathbf{u} = \lambda \mathbf{v} + \mu \mathbf{w}.$$

26 Find all vectors \mathbf{x} that satisfy the equation

$$(-\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}) \times \mathbf{x} = \mathbf{i} + 5\mathbf{j} - 3\mathbf{k}.$$

28. What condition must be satisfied by the nonzero vectors \mathbf{a} and \mathbf{b} to guarantee that the equation $\mathbf{a} \times \mathbf{x} = \mathbf{b}$ has a solution for \mathbf{x} ? Is the solution unique?

Ex. 10.4

8 Find equation of the plane satisfying the given conditions. Passing through the line of intersection of the planes $2x + 3y - z = 0$ and $x - 4y + 2z = -5$, and passing through the point $(-2, 0, -1)$.

18 Find equations of the line specified in vector and scalar parametric forms and in standard form. Through $(2, -1, -1)$ and parallel to each of the two planes $x + y = 0$ and $x - y + 2z = 0$.

28 Find the distance from the origin to the line $x + y + z = 0$, $2x - y - 5z = 1$.

Lecture Note [Ex 3.15](#) (p13),

Exercises for students -[Qu 6](#) (p29)

Homework 2

(Total: 12 questions)

Ex. 11.1

12 Find the velocity, speed and acceleration at time t of the particle whose position is $\mathbf{r}(t)$. Describe the path of the particle.

$$\mathbf{r} = at \cos \omega t \mathbf{i} + at \sin \omega t \mathbf{j} + b \ln t \mathbf{k}$$

24 If the position and velocity vectors of a moving particle are always perpendicular show the path of the particle lies on a sphere.

29 Write the Product Rule for $\frac{d}{dt}(\mathbf{u} \times (\mathbf{v} \times \mathbf{w}))$.

32 Expand and simplify: $\frac{d}{dt}((\mathbf{u} \times \mathbf{u}') \cdot (\mathbf{u}' \times \mathbf{u}''))$.

Ex. 11.3

In exercises 2 and 4, find the required parametrization of the first quadrant part of the circular arc $x^2 + y^2 = a^2$.

2 In terms of the x -coordinate, oriented clockwise.

4 In terms of arc length measured from $(0, a)$, oriented clockwise.

16 Describe the parametric curve \mathcal{C} given by

$$x = a \cos t \sin t, \quad y = a \sin^2 t, \quad z = bt.$$

What is the length of \mathcal{C} between $t = 0$ and $t = T > 0$?

18 Describe the intersection of the sphere $x^2 + y^2 + z^2 = 1$ and the elliptic cylinder $x^2 + 2z^2 = 1$. Find the total length of this intersection curve.

27 Let $\mathbf{r} = \mathbf{r}_1(t)$, ($a \leq t \leq b$) and $\mathbf{r} = \mathbf{r}_2(u)$, ($c \leq u \leq d$), be two parametrizations of the same curve C , each one-to-one on its domain and each giving C the same orientation (so that $\mathbf{r}_1(a) = \mathbf{r}_2(c)$ and $\mathbf{r}_1(b) = \mathbf{r}_2(d)$). Then for each t in $[a, b]$ there is a unique $u = u(t)$ such that $\mathbf{r}_2(u(t)) = \mathbf{r}_1(t)$. Show that

$$\int_a^b \left\| \frac{d}{dt} \mathbf{r}_1(t) \right\| dt = \int_c^d \left\| \frac{d}{du} \mathbf{r}_2(u) \right\| du,$$

and thus that the length of C is independent of parametrization.

Extra questions

1 Verify the formula for the arc length element in cylindrical coordinates,

$$ds = \sqrt{\left(\frac{dr}{dt}\right)^2 + (r(t))^2 \left(\frac{d\theta}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt,$$

given in this section.

2 Verify the formula for the arc length element in spherical coordinates,

$$ds = \sqrt{\left(\frac{d\rho}{dt}\right)^2 + (\rho(t))^2 \left(\frac{d\phi}{dt}\right)^2 + (\rho(t) \sin \phi(t))^2 \left(\frac{d\theta}{dt}\right)^2} dt,$$

given in this section.

3 Reparametrize the curve $\mathbf{r} = a \cos^3 t \mathbf{i} + a \sin^3 t \mathbf{j} + b \cos 2t \mathbf{k}$ ($0 \leq t \leq \pi/2$) in the same orientation in terms of the arc length measured from the point when $t = 0$.