

Moral A PDE has arbitrary functions in its solution. In these examples the arbitrary functions are functions of one variable that combine to produce a function $u(x, y)$ of two variables which is only partly arbitrary.

A function of two variables contains *immensely* more information than a function of only one variable. Geometrically, it is obvious that a surface $\{u = f(x, y)\}$, the graph of a function of two variables, is a much more complicated object than a curve $\{u = f(x)\}$, the graph of a function of one variable.

To illustrate this, we can ask how a computer would record a function $u = f(x)$. Suppose that we choose 100 points to describe it using equally spaced values of x : $x_1, x_2, x_3, \dots, x_{100}$. We could write them down in a column, and next to each x_j we could write the corresponding value $u_j = f(x_j)$. Now how about a function $u = f(x, y)$? Suppose that we choose 100 equally spaced values of x and also of y : $x_1, x_2, x_3, \dots, x_{100}$ and $y_1, y_2, y_3, \dots, y_{100}$. Each pair x_i, y_j provides a value $u_{ij} = f(x_i, y_j)$, so there will be $100^2 = 10,000$ lines of the form

$$x_i \quad y_j \quad u_{ij}$$

required to describe the function! (If we had a prearranged system, we would need to record only the values u_{ij} .) A function of three variables described discretely by 100 values in each variable would require a million numbers!

To understand this book what do you have to know from calculus? Certainly all the basic facts about partial derivatives and multiple integrals. For a brief discussion of such topics, see the Appendix. Here are a few things to keep in mind, some of which may be new to you.

1. Derivatives are *local*. For instance, to calculate the derivative $(\partial u / \partial x)(x_0, t_0)$ at a particular point, you need to know just the values of $u(x, t_0)$ for x near x_0 , since the derivative is the limit as $x \rightarrow x_0$.
2. Mixed derivatives are equal: $u_{xy} = u_{yx}$. (We assume throughout this book, unless stated otherwise, that all derivatives exist and are continuous.)
3. The chain rule is used frequently in PDEs; for instance,

$$\frac{\partial}{\partial x}[f(g(x, t))] = f'(g(x, t)) \cdot \frac{\partial g}{\partial x}(x, t).$$

4. For the integrals of derivatives, the reader should learn or review Green's theorem and the divergence theorem. (See the end of Section A.3 in the Appendix.)
5. Derivatives of integrals like $I(t) = \int_{a(t)}^{b(t)} f(x, t) dx$ (see Section A.3).
6. Jacobians (change of variable in a double integral) (see Section A.1).
7. Infinite series of functions and their differentiation (see Section A.2).
8. Directional derivatives (see Section A.1).
9. We'll often reduce PDEs to ODEs, so we must know how to solve simple ODEs. But we won't need to know anything about tricky ODEs.

EXERCISES

1. Verify the linearity and nonlinearity of the eight examples of PDEs given in the text, by checking whether or not equations (3) are valid.
2. Which of the following operators are linear?
 - (a) $\mathcal{L}u = u_x + xu_y$
 - (b) $\mathcal{L}u = u_x + uu_y$
 - (c) $\mathcal{L}u = u_x + u_y^2$
 - (d) $\mathcal{L}u = u_x + u_y + 1$
 - (e) $\mathcal{L}u = \sqrt{1+x^2}(\cos y)u_x + u_{yxy} - [\arctan(x/y)]u$
3. For each of the following equations, state the order and whether it is nonlinear, linear inhomogeneous, or linear homogeneous; provide reasons.
 - (a) $u_t - u_{xx} + 1 = 0$
 - (b) $u_t - u_{xx} + xu = 0$
 - (c) $u_t - u_{xxt} + uu_x = 0$
 - (d) $u_{tt} - u_{xx} + x^2 = 0$
 - (e) $iu_t - u_{xx} + u/x = 0$
 - (f) $u_x(1+u_x^2)^{-1/2} + u_y(1+u_y^2)^{-1/2} = 0$
 - (g) $u_x + e^y u_y = 0$
 - (h) $u_t + u_{xxxx} + \sqrt{1+u} = 0$
4. Show that the difference of two solutions of an inhomogeneous linear equation $\mathcal{L}u = g$ with the same g is a solution of the homogeneous equation $\mathcal{L}u = 0$.
5. Which of the following collections of 3-vectors $[a, b, c]$ are vector spaces? Provide reasons.
 - (a) The vectors with $b = 0$.
 - (b) The vectors with $b = 1$.
 - (c) The vectors with $ab = 0$.
 - (d) All the linear combinations of the two vectors $[1, 1, 0]$ and $[2, 0, 1]$.
 - (e) All the vectors such that $c - a = 2b$.
6. Are the three vectors $[1, 2, 3]$, $[-2, 0, 1]$, and $[1, 10, 17]$ linearly dependent or independent? Do they span all vectors or not?
7. Are the functions $1 + x$, $1 - x$, and $1 + x + x^2$ linearly dependent or independent? Why?
8. Find a vector that, together with the vectors $[1, 1, 1]$ and $[1, 2, 1]$, forms a basis of \mathbb{R}^3 .
9. Show that the functions $(c_1 + c_2 \sin^2 x + c_3 \cos^2 x)$ form a vector space. Find a basis of it. What is its dimension?
10. Show that the solutions of the differential equation $u''' - 3u'' + 4u = 0$ form a vector space. Find a basis of it.
11. Verify that $u(x, y) = f(x)g(y)$ is a solution of the PDE $uu_{xy} = u_x u_y$ for all pairs of (differentiable) functions f and g of one variable.